

VLT/NACO near-infrared imaging and spectroscopy of N88A in the SMC [★]

G. Testor^{1**}, J.L. Lemaire^{1,2**}, M. Heydari-Malayeri¹, L. E. Kristensen³, S. Diana², D. Field⁴

¹ LERMA, UMR 8112 du CNRS, Observatoire de Paris, 92195 Meudon, France

e-mail: gerard.testor@obspm.fr

² Université de Cergy-Pontoise, 95031 Cergy Cedex, France

³ Leiden Observatory, Leiden University, Niels Bohrweg 2, 2333 CA Leiden, The Netherlands

⁴ Department of Physics and Astronomy, Århus University, 8000 Århus C, Denmark.

Received ...; accepted ...

ABSTRACT

Aims. We present near-infrared imaging and spectroscopic high spatial resolution observations of the SMC region N88 containing the bright, excited, extincted and compact H II region N88A of size ~ 1 pc.

Methods. To investigate its stellar content and reddening, N88 was observed using spectroscopy and imagery in the JHKs- and L'-band at a spatial resolution of ~ 0.1 - $0.3''$, using the VLT UT4 equipped with the NAOS adaptive optics system. In order to attempt to establish if the origin of the infra-red (IR) excess is due to bright nebulosity, circumstellar material and/or local dust, we used Ks vs J-K colour-magnitude (CM) and JHK colour-colour (CC) diagrams, as well as L' imagery.

Results. Our IR-data reveal in the N88 area an IR-excess fraction of ≥ 30 per cent of the detected stars, as well as an unprecedentedly detailed morphology of N88A. It consists of an embedded cluster of $\sim 3.5''$ (~ 1 pc) in diameter, of at least thirteen resolved stars superposed with an unusual bright continuum centered on a very bright star. The four brightest stars in this cluster lie red-ward of H-K ≥ 0.45 mag, and could be classified as young stellar object (YSO) candidates. Four other probable YSO candidates are also detected in N88 along a south-north bow-shaped thin H₂ filament at $\sim 7''$ east of the young central bright star. This star, that we assume to be the main exciting source, could also be complex. At $0.2''$ east of this star, a heavily embedded core is detected in the L'-band. This core with L' ~ 14 mag and L'-K ≥ 4.5 mag could be a massive class I protostar candidate. The $2.12 \mu\text{m}$ H₂ image of N88A resembles a shell of diameter $\sim 3''$ (~ 0.9 pc) centered on the bright star. This shell is formed of three bright components, of which the brightest one superposes the ionization front. The line ratios of H₂ 2-1 S(1) and 1-0 S(0) relative to 1-0 S(1), as well as the presence of high v lines, are indicative of photodissociation regions, rather than shocks.

Key words. galaxies: Small Magellanic Cloud – ISM: individual objects: N88A – Stars: formation – Stars: individual: pre-main sequence

1. Introduction

The Small Magellanic Cloud (SMC) is rich in H II regions and young OB associations. Because of its known and relatively small distance (~ 65 kpc) (Kovacs 2000), its face-on position relatively free from foreground extinction (McNamara & Feltz 1980), and low internal extinction (Westerlund 1997), it is well suited to study both individual stars and very compact objects, as well as global structures. It is an ideal laboratory for studying the formation and evolution of massive stars in a low metallicity environment. Understanding the characteristics of massive stars and their interaction with their environment is a key problem in astrophysics. We have made some progress concerning the early stages of massive star formation in the galaxy, but the current knowledge about the early stages of massive star evolution in other galaxies is mediocre at

best. The main reason is that the earliest stages of massive star evolution are deeply enshrouded, inaccessible in the optical wavelengths. Another reason is that these stars are often members of very crowded regions. Today, high-spatial near-infra-red (NIR) resolution observations using NACO attached to the VLT, are able to overcome these obstacles in the SMC. Our search for the youngest massive stars in the Magellanic Clouds (MCs) (Heydari-Malayeri & Testor 1982) led to the discovery of a distinct and very rare class of H II regions that we called high-excitation compact H II "blobs" (hereafter HEBs) listed in Testor (2001). In contrast to the ordinary H II regions of the MCs, which are extended structures spanning several arcminutes on the sky (> 50 pc) and are powered by a large number of hot stars, HEBs are very dense small regions, usually $2''$ to $8''$ in diameter (0.8 to 3 pc), ionized by one or a few massive stars and affected by local dust. Two other HII regions, MA 1796 and MG 2 (less than 1pc across) heavily extincted and ionized by a small young cluster, have been found by Stanghellini et al. (2003) in the SMC.

In the present paper we focus on the peculiar HEB LHA 115-N88A, hereafter labelled N88A of diameter ~ 1 pc

Send offprint requests to: gerard.testor@obspm.fr and jean-louis.lemaire@obspm.fr

[★] Based on observations obtained at the European Southern Observatory, El Paranal, Chile

^{**} visiting astronomer at VLT Paranal

(Testor & Pakull 1985) in the extended H II region LHA 115-N88 or N88 (Henize 1956) of diameter ~ 10 pc. N88 lies in the Shapley Wing of the SMC and contains the young cluster HW 81 ($\sim 0.6'$) (Hodge & Wright 1977). It is known that the SMC is made of four H I layers with different velocities (McGee & Newton 1982). This situation complicates the study, particularly in the region of the Shapley wing. However, the available H I observations provide helpful data for the study of this region. In particular, the N88 region lying at about $35'$ (~ 700 pc) west of N83/N84 is not apparently associated with the H I cloud of these H II regions (Heydari-Malayeri et al. 1988). N88A should be associated with the H I component of velocity $+134 \text{ km s}^{-1}$ (McGee & Newton 1982).

N88A is the brightest and the most excited HEB in the MCs. It is also the most reddened H II region in these galaxies of low-metallicity (Heydari-Malayeri et al. 2007). Numerous optical detailed studies have been made on this object (Testor & Pakull 1985, Wilcot 1994, Heydari-Malayeri et al. 1999 (hereafter HM99), Kurt et al. 1999, Testor et al. 2003). Nevertheless, many uncertainties remain to understand the true nature of N88A, such as its exciting source, that still remains unidentified, as well as the nature of the reddening.

Israel & Koornneef (1988, 1991) have detected the presence of molecular hydrogen in N88 through H_2 emission which is either shock-excited on a small scale of $0.46''$ by stars embedded in the molecular cloud, or radiatively excited on a large scale ($3''$ - $60''$). However, their low spectral ($R=50$) and spatial (7.5 - $10''$ aperture) resolutions did not allow discrimination of these different processes. They described N88A as a strong NIR source dominated by nebular emission containing a strong hot dust component and noticed that N88A has an unusual blue J-H colour. In Testor et al. (2005), at low spatial resolution, a pure H_2 emission is detected in N88A as well as along a south-north diffuse long filament at ~ 6 - $8''$ to the east. In N66 (Henize 1956) a giant HII region in the SMC, Schmeja et al. (2009) have reported that most of the H_2 emission peaks coincide with the bright component of the ionized gas and with compact embedded young clusters where candidate YSOs have been identified. Using SEST, Israel et al. (2003) detected a CO molecular cloud of $1.5' \times 1.5'$ in the region, reporting spectra and maps of the ^{12}CO lines J=1-0 and J=2-1. Stanimirovic et al. (2000) found that the highest values of the dust-to-gas mass ratio and dust temperature in the SMC are found in N88A.

IR studies of a similar size young star formation region like the Trapezium region in Orion ($\sim 0.75 \times 0.75$ pc) (Lada et al. 2000) and the more extended 30 Doradus in the LMC (Maercker & Burton 2005) showed that during star formation, YSOs are associated with the circumstellar material inducing IR-excess emission, and also that the use of JHK CC diagrams are useful tools to detect this emission. However, for young massive stars generally found in embedded clusters, their surrounding material destruction time scale is short, making their observation difficult (Bik et al. 2005, 2006). The most suitable wavelength to determine the nature of the IR-excess is the L-band, increasing the IR-excess and reducing the contribution of extended emission from reflection nebulae and H II regions (Lada et al. 2000). IR-excess can be useful to determine the origin of the reddening in embedded young clusters. Martin-Hernandez et al. (2008) found in N88A, from a mid-IR high spatial res-

Table 1. Log of the photometric VLT/NACO observations.

Id.	Filter	Expo. t(s) x n	Mode	Date	Seeing FWHM	
					($''$)	($''$)
N88A	J $1.27 \mu\text{m}$	20 x 30	-	-	-	0.35
	H $1.66 \mu\text{m}$	20 x 30	-	-	-	0.27
	Ks $2.18 \mu\text{m}$	30 x 30	S54	9/10/2004	0.6-0.9	0.21
	L' $3.8 \mu\text{m}$	0.2 x 26	S27	10/10/2004	0.9-1.2	0.10
	NB-2.12	200x 20	S54	-	-	0.19
	NB-2.24	150x 20	S54	-	-	0.25
	Ks	60 x 30	S27	11/10/2004	-	0.10

Table 2. Log of the VLT/NACO long-slit spectroscopic observations.

Id.	Date	Slit (mas)	Exposures t(s) x n	Mode	$\lambda/\delta\lambda$ Seeing	
					($''$)	($''$)
S1	9/10/2004	172	100 x 10	S54-4-SHK	400	0.6-0.9
S2	10/10/2004	-	-	-	-	-
S3	11/10/2004	-	-	-	-	-
hip8485	9/10/2004	-	2.5 x 4	-	-	-
hip29968	9/10/2004	-	2 x 4	-	-	-
hip103087	11/10/2004	-	1.8 x 4	-	-	-

olution Spitzer-IRS spectrum, a rising dust continuum and PAH bands, typical characteristics in H II regions. Using radio observations, Indebetouw et al. (2004) found that N88A is ionized by an O5 type star.

In the present paper we present the results of JHK- and L'-band high spatial resolution observations of N88A and its surroundings, using adaptive optics at the VLT. Section 2 discusses the instrumentation employed during these observations, and the data acquisition and reduction procedures used. Section 3 describes the results and analysis of our observations, and Section 4 gives our conclusions.

2. Observations and data reduction

NIR observations of N88A were obtained at the ESO Very Large Telescope (VLT) during October 2004. Images and spectra were taken using NACO on UT4, composed of the Nasmyth Adaptive Optics System (NAOS) and the High Resolution IR Camera and Spectrometer (CONICA). The detector was a 1026×1024 SBRC InSb Alladin 3 array. The cameras S54 and S27 were used in the range 1.0 - $2.5 \mu\text{m}$ and the L27 camera in the range 2.5 - $5.0 \mu\text{m}$.

The field-of-view (FOV) of the S54 camera was $54'' \times 54''$ with a pixel size of $0.05274''$, corresponding to 0.015 pc at the distance modulus of 19.05 for the SMC (Kovacs 2000). The FOV of the S27 and L27 camera was $27.15'' \times 27.15''$ with a pixel size of $0.02637''$, corresponding to 0.0075 pc. For spectroscopy we used the S54 camera.

As adaptive optics (AO) reference source for wavefront sensing we used the object itself (N88A) (Testor et al. 2005). The conditions were photometric and the seeing ranging from $0.65''$ to $1''$ in the visible. After subtraction of the average dark frame, each image was divided by the normalized flat field image. The data were reduced mainly with the ESO software packages MIDAS and ECLIPSE.




Fig. 1. JHK colour composite image of SMC N88 containing the bright HEB N88A. The image size extracted from the S54 camera field corresponds to $50'' \times 50''$ (or $\sim 14 \text{ pc} \times 14 \text{ pc}$). The lower spatial density of faint stars close to the edges of the frame delimits approximatively the cluster N88-cl. In the north-west quadrant the lack of stars is explained by the presence of the molecular cloud (HM99). A red filament is visible east of N88A. J, H and K are respectively coded B, G and R. North is up and East is left.

2.1. Imaging

Images through J, H, Ks broad-band and $2.12\mu\text{m}$, $2.24\mu\text{m}$ narrow band filters were obtained with the S54 camera. A composite JHKs colour image of the observed field is shown in Fig. 1. Images with higher spatial resolution, in the Ks and L' large band, were also obtained with the S27 camera. The log of NIR imaging observations is given in Table 1. The AutoJitter mode was used: that is, at each exposure, the telescope moves according to a random pattern in a $6'' \times 6''$ box. Table 1 lists the stellar full-width-at-half-maximum (FWHM) in final images of different observed bands of the star at J2000 coordinates $(\alpha, \delta) = (1^{\text{h}}, 24^{\text{m}}, 8.88^{\text{s}}, -73^{\circ}, 8', 56.2'')$ of the 2MASS survey (Cutri et al. 2003). The AO image is affected by anisoplanatism and leads to degradation of the point spread function (PSF) becoming more elongated as the angular offset from the guide star increases. This has been taken into consideration for the photometric measurements, as explained in Sect. 2.3.

2.2. Spectroscopy

Spectroscopy was performed in the S54-4-SHK mode (broad-band filter), giving a linear dispersion of 1.94 nm/pixel and a spatial scale of 53 mas/pixel . Three long-slit spectra S1, S2, and S3 were chosen from the IR images given by NACO. S1 ($\text{PA} = 115^{\circ}$) and S2 ($\text{PA} = 89.3^{\circ}$) cross the central bright star. S3 ($\text{PA} = 18.9^{\circ}$) crosses the stars #37 and #47 (see Fig. 3). The slit width was 172 mas and the spectral resolution ~ 400 . For each exposure the detector integration time (DIT) was 100s . Ten exposures were obtained in the Autonod on slit mode, which allows spectroscopy of moderately extended objects. The log of spectroscopic observations is given in Table 2. In order to remove telluric absorption features, stars with a similar airmass were observed as telluric standards. Spectroscopy was reduced with the MIDAS software package LONG.




Fig. 2. Finding chart in the J-band obtained with the S54 camera (Field 1). The inset corresponds to an enlargement (factor 2) and unsaturated image of N88A extracted from Field 1. The numbering refers to Table 3. The total field size corresponds to $48.6'' \times 47.5''$ (or $\sim 14 \text{ pc} \times 14 \text{ pc}$).

2.3. Photometry

In Fig. 2 we present the N88 region observed with the S54 camera (Field 1) through the J-band filter. The instrumental magnitudes of the elongated stars (see Sect. 2.1) outside the central region, were derived with DAOPHOT (Stetson 1987), using concentric aperture photometry to integrate all the flux of each star. Although PSF photometry is better adapted for crowded fields, we could not use it. Indeed the stars in our field were too faint and/or crowded to obtain the number of PSF stars necessary to use the photometric analysis elaborated by Pugliese et al (2002) taking into account the AO anisoplanatism effect.

The detected stars are identified by a number referring to Column 1 of Table 3 that gives the astrometry and photometry. The central object N88A is not affected by anisoplanatism, so the JHK instrumental magnitudes of the stars were derived using the DAOPHOT's multiple-simultaneous-profile-fitting photometry routine (NSTAR), well adapted for photometry in crowded fields. The detected stars are shown in the inset of Fig. 2 and are also listed in

Table 3. Almost all the stars of Field 1 (Fig. 2) have photometric uncertainties in the J-, H- and K-band, less than 0.03 mag for stars with $K < 16$ mag, less than 0.06 for stars with $16 < K < 18$ mag and greater than 0.1 for stars $18 < K < 20$ mag.

Fig. 3 (Field 2) shows only a part of N88 observed with the S27 camera through the Ks-band filter. On this figure the directions of the spectra S1, S2 and S3 are plotted. This camera, with a pixel two times smaller than S54, has a better spatial resolution, so the analysis of the crowded field of N88A with the routine NSTAR allows a more accurate star detection. In Fig. 4a the stars of N88A are identified by a number referring to Column 1 of Table 4. In this table, the average photometric errors of the stars reported by DAOPHOT are ~ 0.04 mag in J and 0.06 in H and Ks for stars of magnitude ≤ 16 except for the bright isolated star #57 outside N88A (~ 0.03 and ~ 0.05 mag). This star will be used as PSF. The photometric calibration was obtained using the isolated 2MASS star at J2000 (α, δ) = ($1^h, 24^m, 8.88^s$, $-73^\circ, 8', 56.2''$) corresponding to our star




Fig. 3. Ks-band image of SMC-N88 obtained with the S27 camera (Field 2). The location of the slits labelled S1, S2 and S3 used in the spectroscopic mode is indicated by a solid line. In N88A the positions of the stars #37, #41, #42 and #47 are indicated. The bottom left is an enlargement of $0.55'' \times 0.55''$ containing the complex object N88B. The image size corresponds to $25'' \times 25''$ (or $\sim 7 \text{ pc} \times 7 \text{ pc}$).

#57. The conversion of pixel coordinates to α and δ was derived using the same star and the relative positions of our stars are accurate to better than $0.1''$.

In the core of N88A the determination of the sky aperture parameters used in NSTAR is very sensitive, even with the S27 camera. Indeed, the wings of the stars superpose the wings of the strong continuum. The distribution of this continuum resembles a gaussian profile (Fig. 5). The error on the magnitude of these stars, due to a steeply sloping continuum background, is greater than the error given by DAOPHOT. Because of this situation, the K magnitude of the central star labelled #41 at low and at higher spatial resolution (Field 1 and Field 2) is respectively 13.82 and 14.99 mag. Therefore, the K magnitudes of the stars #37, #41, and #47 (Field 2) were remeasured by subtracting a one-dimensional (1-D) profile corresponding to the PSF crossing the center of the isolated reference star # 57. The magnitude of the PSF was multiplied by a factor in such a way that only the continuum remains visible. In this case

its magnitude corresponds to the magnitude of the star. An example is given in Fig. 5. The K magnitudes of the stars obtained with this method are listed in col. 7 of Table 4. In this table the magnitude of #41 is in agreement with the magnitude obtained with NSTAR (col 5), while for #37, #42 and #47 the Ks magnitude is greater. Several faint stars under the detection level, or slightly extended, are not detected or rejected by DAOPHOT (Fig. 4b). In our underexposed L'-band frame of N88, not shown here, contrary to the JHK-band, no star is found except star #57. However, in N88A the bright star #41 is visible as well as the stars #37, #42, #47 and several unresolved features. A peculiar bright core labelled L1-C (see Sect. 3. below) located $0.2''$ east of #41 is also found. This core coincides approximatively with the HST absorption lane (HM99) and has a very faint counterpart in the Ks-band (Fig. 6). In order to derive the L' magnitude of these objects, we referred to the L' photometry of Israel & Koorneef (1991), obtained through a $7''$ aperture (Table 5). The integration of N88A




Fig. 4. Enlargements of the HEB N88A in Field 2. a) Ks-band obtained with the S27 camera. The numbering refers to Table 4 (Field 2). b) the DDP process applied to the Ks-band allows to enhance the faint stars of N88A-cl. A few stars not identified with DAOPHOT and small features are indicated by arrows. c) Stromgren y image obtained with the HST showing the absorption lane, the stars #1, #2 as well as the ‘dark spot’ of HM99. The Stars #1 and #2 correspond to our stars #41 and #42. d) L'-band obtained with the S27 camera, the different components are indicated. The field corresponds to $4.43'' \times 4.32''$ (or $1.27 \text{ pc} \times 1.27 \text{ pc}$) and is outlined in (Fig. 3).

in our sky subtracted Ks image, using a $4''$ aperture, gives a magnitude of 11.08, in agreement with Israel & Koorneef (1991) and other authors (Table 5). In an aperture of $4''$ we have integrated the L' flux of N88A that was then calibrated with $L' = 8.92 \text{ mag}$ given by Israel & Koorneef (1991). The L' magnitudes of the stars #57, #41 and the core L1-C (continuum subtracted) were obtained using an aperture of $0.35''$ and are listed in Table 4.

3. Results and discussion

3.1. The HEB N88A

The HST $H\alpha$ (F656N) and continuum Stromgren y (F547M) images of N88A described in HM99, show two inhomogeneous wings separated by a north-south absorption lane (Fig. 4c). The western wing is much brighter and contains two faint stars, #1 and #2, corresponding to our

stars #41 and #42, as well as a ‘dark spot’ to the south at the location of our star #37. Fig. 7 shows the intensity distribution plots in the Stromgren y- and Ks-band in the direction of the slit S1 (Fig. 3) crossing N88B and the star #41. In Fig. 7 the stars #41 and #42 of magnitude 14.99 and 16.11 respectively (Table 4) coincide with the two faint stars #1 and #2 (Fig. 4c) of $y = 18.2 \text{ mag}$ and 18.3 mag (HM99).

Through the Ks filter (Figs. 4a and b), N88A appears as a circular nebular region of $\sim 3.4''$ diameter centered on the relatively bright star #41. This star coincides with the 2MASS point source 012407.92-730904.1 of $K = 11.18 \text{ mag}$ (Cutri et al. 2003). In a diameter of $\sim 3.6''$ our N88A image exhibits a small embedded cluster labelled N88A-cl of at least thirteen stars (Fig. 4b). In Figure 4b the usual Digital Development Process (DDP) introduced by Okano was applied to enhance the faint stars by compressing the range

Table 3. Stars in the region of N88A observed with the S54 camera (Field 1).

Id.	$\alpha(2000)$	$\delta(2000)$	mag-J	mag-H	Mag-K	J-K	J-H	H-K
1	1 24 11.42	73 9 26.15	18.24	17.65	17.50	0.74	0.59	0.15
2	1 24 11.86	73 9 25.85	17.63	17.71	17.65	-0.02	-0.08	0.06
3	1 24 8.67	73 9 23.08	19.08	18.88	18.84	0.24	0.20	0.04
4	1 24 13.84	73 9 22.77	18.46	18.48	18.45	0.01	-0.02	0.03
5	1 24 7.69	73 9 20.63	17.89	17.61	17.34	0.55	0.28	0.27
6	1 24 11.02	73 9 20.11	15.76	15.85	15.82	-0.06	-0.09	0.03
7	1 24 6.85	73 9 19.56	19.16	18.71	18.51	0.66	0.45	0.21
8	1 24 13.75	73 9 18.85	15.17	15.37	15.38	-0.21	-0.20	-0.01
9	1 24 8.72	73 9 18.79	18.22	18.18	18.02	0.20	0.04	0.16
10	1 24 7.03	73 9 18.05	18.80	18.79	18.71	0.09	0.01	0.08
11	1 24 12.53	73 9 15.61	16.27	16.45	16.47	-0.20	-0.18	-0.02
12	1 24 7.30	73 9 15.25	16.72	16.87	16.87	-0.16	-0.15	-0.00
13	1 24 10.75	73 9 14.84	18.67	18.53	18.06	0.61	0.14	0.47
14	1 24 5.18	73 9 14.70	16.27	16.44	16.31	-0.04	-0.17	0.13
15	1 24 7.45	73 9 14.32	17.31	17.48	17.47	-0.15	-0.17	0.01
16	1 24 5.16	73 9 14.07	16.19	16.40	16.26	-0.07	-0.21	0.14
17	1 24 12.57	73 9 13.33	17.44	17.59	17.40	0.04	-0.15	0.19
18	1 24 8.97	73 9 11.82	19.78	19.65	19.52	0.26	0.12	0.14
19	1 24 9.30	73 9 10.17	18.44	18.21	17.87	0.57	0.23	0.34
20	1 24 9.13	73 9 10.09	18.30	18.23	18.19	0.11	0.08	0.03
21	1 24 9.58	73 9 9.84	19.11	18.51	17.66	1.45	0.60	0.84
22	1 24 5.76	73 9 9.34	18.89	18.59	17.93	0.96	0.30	0.66
23	1 24 8.58	73 9 9.01	19.65	19.34	18.95	0.70	0.31	0.39
24	1 24 9.93	73 9 8.68	18.66	18.36	17.80	0.86	0.31	0.55
25	1 24 11.52	73 9 8.03	18.89	18.73	18.51	0.38	0.15	0.22
26	1 24 10.13	73 9 7.70	19.48	19.46	19.30	0.17	0.02	0.15
27	1 24 9.45	73 9 7.50	18.37	18.18	17.97	0.40	0.19	0.20
28	1 24 4.96	73 9 7.42	16.32	16.48	16.42	-0.10	-0.16	0.06
29	1 24 9.25	73 9 7.17	18.21	18.11	17.88	0.33	0.10	0.24
30	1 24 6.14	73 9 6.35	17.85	17.51	17.33	0.52	0.34	0.18
31	1 24 9.09	73 9 5.99	15.74	15.73	15.51	0.23	0.01	0.16
32	1 24 7.57	73 9 5.80	19.04	19.09	18.91	0.13	-0.05	0.18
33	1 24 9.98	73 9 5.77	19.46	18.76	17.83	1.63	0.70	0.93
36	1 24 5.82	73 9 5.09	17.36	17.43	17.26	0.10	-0.07	0.17
37 ⁿ	1 24 7.84	73 9 4.24	15.01	15.10	14.28	0.73	-0.09	0.81
41 ⁿ	1 24 7.96	73 9 3.74	14.68	14.56	13.82	0.86	0.12	0.74
42 ⁿ	1 24 7.85	73 9 3.60	15.05	15.05	14.44	0.61	0.00	0.61
47 ⁿ	1 24 7.96	73 9 3.00	15.55	15.36	14.88	0.67	0.19	0.48
46	1 24 10.08	73 9 3.05	19.65	18.93	18.18	1.47	0.72	0.75
48	1 24 11.02	73 9 2.86	18.93	18.90	18.82	0.12	0.03	0.09
50 ⁿ	1 24 7.53	73 9 2.09	17.61	17.62	17.16	0.45	-0.01	0.46
52	1 24 7.01	73 9 1.79	18.80	18.98	18.57	0.23	-0.18	0.41
53	1 24 10.35	73 9 0.80	19.91	19.51	18.95	0.96	0.40	0.56
54	1 24 12.17	73 8 59.81	17.75	17.59	17.29	0.46	0.16	0.30
55	1 24 8.53	73 8 57.45	20.05	20.01	19.65	0.40	0.04	0.36
56	1 24 9.27	73 8 56.87	17.57	17.42	17.29	0.28	0.15	0.13
57	1 24 9.07	73 8 56.24	15.36	14.88	14.59	0.77	0.48	0.29
58	1 24 10.81	73 8 55.78	18.45	18.43	18.26	0.19	0.02	0.17
59	1 24 9.86	73 8 54.07	16.51	16.67	16.64	-0.13	-0.16	0.03
60	1 24 3.68	73 8 53.33	18.51	18.03	17.59	0.92	0.48	0.44
61	1 24 8.36	73 8 53.17	19.27	18.65	18.44	0.83	0.62	0.21
62	1 24 3.59	73 8 50.89	18.06	17.65	17.43	0.63	0.41	0.22

ⁿ Stars in N88A analyzed using the NSTAR routine.

of brightnesses between the bright and dim portions of the image. The K photometry of these stars is listed in Table 4. These stars mainly concentrated to the north and east, superpose numerous nebular structures. Interestingly these stars are aligned in the direction of the interface between the HII regions N88A and N88B (HM99). Through the L'-band, N88A seems to be essentially formed by four distinct components labelled L1, L2, L3 and L4 (Fig. 4d). L1, that contains the core labelled L1-C (Fig. 7), is the brightest and most compact component. In the Ks-band L1-C (Figs. 4a and 6) shows a very faint counterpart ($L-K_s \geq 4.2$). L2 and L3 are more diffuse and coincide with the stars #47 and #37. L4 is bright, extended and formed by two east-west elongated subcomponents spanning between stars #47 and #41. In the L'-band the star #41 is relatively bright ($L' \sim 14.1$ mag) and well resolved (Fig. 4d). All these components superpose a diffuse nebular continuum. On the y (F547M) continuum image (Fig. 4c) the 'dark spot' corresponding to our component L3 is very bright. The y continuum structures located between stars #41 and #37, as well as north

to the 'dark spot' (Fig. 4c), are not seen in the L'-band (Fig. 4d).

3.2. The N88B region

At the optical wavelengths, HM99 found that the central star of N88B, corresponding to our star #31, has an integrated magnitude of $y = 16.57$ and consists of at least three components. Our high spatial resolution Ks-band image also shows that star #31 of magnitude $K = 15.74$ and $J-K = 0.23$ is complex and formed of at least three components visible in the inset of Fig. 3. Two of them oriented approximately south-north are relatively bright, whereas the one to the north-east corresponds to a faint diffuse feature. To the east of N88B lies a red bow-shaped filament with a curvature radius of $\sim 3''$ (Fig. 1) centered on N88B (see Sect. 3.6). This filament coincides with the narrow filament north-east of component B detected in the H α -band (HM99).

Table 4. Stars in N88A observed with the S27 camera (Field 2) in the Ks-band (numbers refer to Fig. 4.

Id.	$\alpha(2000)$	$\delta(2000)$	K^{dao}	err	K^{psf}	L^{ap}	K-L
57	1 24 8.88	-73 8 56.00	14.64	0.003		14.36 $\pm$ 0.05	
34	1 24 7.94	-73 9 5.69	18.45	0.070			
35	1 24 7.64	-73 9 5.42	18.73	0.050			
37	1 24 7.83	-73 9 4.56	16.43	0.032	16.66 $\pm$ 0.15		
38	1 24 8.30	-73 9 3.85	18.46	0.060			
40	1 24 8.21	-73 9 3.80	18.40	0.065			
41	1 24 7.93	-73 9 3.99	14.99	0.010	15.05 $\pm$ 0.10	14.1 $\pm$ 0.10	0.95
42	1 24 7.85	-73 9 3.82	16.11	0.047	16.60 $\pm$ 0.30		
43	1 24 8.21	-73 9 3.30	17.62	0.066			
44	1 24 8.23	-73 9 3.05	17.76	0.060			
45	1 24 8.16	-73 9 3.14	17.12	0.044			
47	1 24 7.93	-73 9 3.19	15.62	0.036	15.99 $\pm$ 0.20		
49	1 24 8.16	-73 9 2.70	17.41	0.061			
51	1 24 7.49	-73 9 2.29	18.27	0.040			
52	1 24 7.88	-73 9 1.90	18.65	0.061			
L1-C	1 24 7.95	-73 9 3.80	≥ 18.5	0.024		13.98 $\pm$ 0.20	≥ 4.52

dao , psf and ap magnitudes derived using DAOPHOT, the PSF of star #57, and an aperture respectively.




Fig. 5. The thin solid lines show the profile crossing the center of star #41 without subtraction of the continuum. The dotted line shows the profile (PSF) of the isolated star #40 labelled #80 in Field 1. The intensity of the PSF of FWHM = 0.11'' is multiplied by a normalization factor. The thick solid line shows the continuum obtained after the subtraction of the PSF. The plot range corresponds to 2.64''.

3.3. JHK CM and CC diagrams

Fig. 8 shows a Ks vs. J-K CM diagram of the N88 region. The magnitudes of the stars (Field 1) belonging to N88A are systematically underestimated (see Sect. 2.3). This difference is visible on the diagram where the stars of N88A (Field 2) analysed using the PSF of star #57 (Table 4) are overplotted. However, taking into account the PSF given by DAOPHOT for JHK of about 0.2-0.3'' we assume that the J-K colour values as well as J-H and H-K are correct within the uncertainties. The color excess $E(B-V)$ towards N88 derived for hot stars from the Magellanic Clouds Photometric Survey (Zaritsky et al. 2002) in a radius of 1', is small ~ 0.15 . On Fig. 8 the reddening track for O stars is plotted, assuming a total visual extinction $A_V = 5.8E(J-K)$ (Tapia et al. 2003) and $A_K = 0.112A_V$ (Rieke & Lebosky 1985). It was derived using as reference the star #8 of type O9.5 in




Fig. 6. East-west intensity distributions along the direction of slit S2 extracted from the Ks image (thin line), L' image (thick line) and Stromgren y (HST) images (dotted line). The L'/Ks ratio (dashed line) shows that the core L1-C (see Sec. 3.3.1) coincides with the minimum intensity of the absorption lane on the Stromgren y plot. The plot range corresponds to 5.1''.

Wilcots 1994b. For this star we adopted a $(J-K)_0$ of -0.15 mag (Lejeune & Shaerer 2001).

Several isochrones with different ages corresponding to $Z = 0.004$ are overplotted (Fig. 8). The diagram appears to reveal two populations. The first one is a young population of dwarf and massive O stars which appears to be fitted with the 3 Myr isochrone. The second one could be a clump of red giant stars of K magnitude in the range of 17-19.5 expanding in the age 300 Myr-10 Gyr. The stars lying beyond the 10 Gyr isochrones are likely to represent embedded stars situated deeper in the molecular cloud, young stars with circumstellar material or evolved stars surrounded by dust.

Fig. 9 shows the H-K vs. J-H CC diagram. In this figure the solid line represents the reddening vector up to $A_V = 5$ mag. All stars that lie on the right side of the reddening vector should have IR-excess. Due to uncertainties on H-K colour we take into account only the stars beyond ~ 0.1




Fig. 7. Intensity distributions along the direction of slit S1 extracted from the Ks-band image (red line) and HST Stromgren y images (black line). The Stars #1, #2 and the absorption lane (HM99) as well as our star #41 and #42 are indicated. The plot range corresponds to $10.6''$.




Fig. 8. J-K CM diagram for the point sources measured in N88 and N88A (Field 1). From left to right are overplotted the 1.4 Myr, 3 Myr, 300 Myr, 1 and 10 Gyr isochrones (extinction free). Some masses between $3M_{\odot}$ and $60M_{\odot}$ are marked with a tick on the 3 Myr isochrone. The stars #37, #41, #42 and #47 of Field 2 (Table 4) analyzed using NSTAR (crosses) and the PSF of star #57 (filled squares) are overplotted.

mag to the right of the reddening vector. Hence, the number of stars with IR-excess extracted from the CC diagram corresponds to at least 30% of the measured stars. On the CC diagram, obtained after integration in an aperture of $4''$ diameter (Table 5), N88A is overplotted and is found to the extreme right (Fig. 9). The plot shows a red J-H colour of 0.33 mag which contrasts with the blue J-H colour given by Israel & Koorneef (1991).




Fig. 9. CC diagram (J-H vs. H-K) for the measured stars in N88 and N88A. The blue line is the main sequence for stars of age 1 Myr and the red line for stars of age 10 Myr. The empty square represent the colour-colour of N88A integrated in an aperture of $4''$ diameter and the filled squares the bright stars in N88A-cl. The solid line is the reddening vector up to $A_V = 5$.

Table 5. Integrated NIR Photometry of the whole region N88A

Reference	J	H	K	L'	J-K	Ks-L'	FWHM	A_V
							($''$)	
Denis ^d	12.19		11.11		1.08		4	7.3
2MASS ^m	12.31	11.98	11.18		1.13		4	7.3
Israel ⁱ			11.05	8.92		2	10	
this paper	12.15	11.92	11.08				4	7.1

^d Cioni et al. (2000), ^m Cutri et al. (2003), ⁱ Israel & Koorneef (1991). From the Infrared Array Camera (IRAC) archive, Charmandaris et al. (2008) using an aperture of $\sim 7''$, give for the wavelength $3.6\mu\text{m}$ (L-band), $4.5\mu\text{m}$, $5.8\mu\text{m}$ and $8\mu\text{m}$ a magnitude for N88A of 9.52, 8.29, 7.1 and 5.58 mag respectively.

3.4. Search for YSO candidates in N88A and N88

Due to their IR-excess emission, YSOs are positioned in the redder parts of the CM and CC diagrams. We first examined low spatial resolution ($\sim 2.5''$) mid-IR Spitzer data of N88A (Charmandaris et al. 2008), and then the near-IR stellar content of both the regions N88A and N88 obtained with the high spatial resolution allowed by NACO ($\sim 0.10 - 0.3''$). On the CM plot $[3.6]-[8]$ versus $[8]$, presented by Charmandaris et al. (2008), N88A lies at the border of the box representing the domain of Class II YSOs. N88A appears also inside the H II region domain. Similarly, on the CC $[5.8]-[8]$ versus $[3.6]-[8]$ diagram N88A is located near the H II region domain, but outside Class I and Class II YSO areas. These observations are explained by the fact that N88A is above all a very bright H II region with strong nebular emission lines and affected by heavy extinc-




Fig. 10. One-dimensional spectra of the stars #37, #41, #42 and #47 plotted in the range of 2.04-2.41 μm .




Fig. 11. One-dimensional nebular spectra plotted in the range of 2.04-2.41 μm .

tion from local dust. In fact the Spitzer data represent flux integrations over the whole H II region ($\sim 1 \text{ pc}^2$). Therefore, detecting a YSO inside the H II region seems hazardous unless the YSO is the dominant source inside N88A, which obviously is not the case. Note that although on the [5.8]-[8] versus [3.6]-[8] colour diagram, based on model calculations (Whitney et al. 2004), N88A appears among Class 0 and Class I data points. This diagram is not applicable to the case of N88A for the reasons explained earlier.

In order to probe the presence of YSOs inside N88A we used our high resolution JHK data. The stars #37, #41,

#42 and #47 in N88A-cl are located at the upper part of the CM diagram (Fig. 8). In the JHK CC diagram (Fig. 9) these stars exhibit an H-K color ranging from 0.48 to 0.81 mag, and can be YSO candidates according to the JHK CC diagram of Maercker & Burton (2005). However, their positions on the J-K versus K diagram (Fig. 8) suggest heavily reddened main-sequence (MS) massive stars of masses between $15M_{\odot}$ and $30M_{\odot}$. Their positions between the 300 Myr and 1 Gyr isochrones are also compatible with supergiants. If confirmed as supergiants, these stars would not be physically associated with N88A, which is very young. The assumption of reddened MS massive stars or massive YSOs seem more plausible. It is very difficult to distinguish between these possibilities. Consequently, caution must be applied, using only JHK band observations to infer circumstellar material fraction in strong nebulous environments. The high spatial, but low spectral resolution S1, S2 and S3 spectra crossing the stars #37, #41, #42 and #47 (see Fig. 10) do not allow to analyze accurately the Br γ line emission profile, which is characteristic for YSO sources, and new spectroscopic observations are needed to clarify the nature of these stars. In N88A, L1 presents a relatively bright peak (L1-C) in the L'-band (Fig. 6). With $K-L' \geq 4.5$, L1-C can be interpreted as a deeply embedded protostar (Lada et al. 2000). L1-C with a FWHM slightly larger than the PSF should still be in its contraction phase, surrounded by a dust shell. With a magnitude of 14 and a strong IR-excess (Table 4) L1-C could be a massive protostar of Class I.

On what concerns the region outside N88A, from our high spatial resolution data the JHK CC diagram (Fig. 9), taking into account the uncertainties excluding some stars close to the reddening vector, we show hereabove that in the extended N88 region, at least 30% of the faint detected stars have an IR-excess. These reddened stars seem to belong to a cluster of faint stars labeled N88cl (Fig. 1) coinciding with the young cluster HW81 (Hodge & Wright 1977) formed of bright stars, situated towards N88 and not affected by dust (HM99). The JHK CC diagram (Fig. 9) shows that the bright stars #6, #8, #11, #12, #15, #28 and #59 in HW81 have no IR-excess and their masses spread in the range of $15M_{\odot}$ to $30M_{\odot}$. In N88-cl most of the reddened stars have a mass $\leq 12M_{\odot}$ (Fig. 8) and are probably intermediate-mass YSO candidates. We assume that N88-cl belongs to N88 (Fig. 1). The stars #21, #33 and #46 of N88-cl aligned on the east filament of N88B (Fig. 1) with an H-K excess > 0.7 mag, could be good YSO candidates (Fig. 9). Their alignment suggests that their formation may be triggered by the expansion of the shell around N88B.

3.5. The ionizing sources

3.5.1. N88A

At 3cm radio emission, Indebetouw et al. (2004) found for N88A a Lyman continuum flux of $\log N_{Lyc} = 49.5$. Using the spectral classification of Smith et al. (2002) we estimate, from this flux, the type of the ionizing source of N88A to range from O4 to O5. The type of the ionizing source derived by HM99 using the H β flux corresponds to an O6 star. The extracted 1-D spectra at different positions along the slits S1, S2 and S3 crossing N88A (Fig. 3) are listed in Table 6. In each position the rows are averaged and the corresponding 1-D spectra are shown in Fig. 10 for the stars and Fig. 11 for nebular emission. The positions of these




Fig. 12. **a)** Relative intensity distributions along slit S1 aligned on stars #31, #41 and #42 (Fig. 2). **b)** Relative intensity distributions along slit S2 crossing the star #41. **c)** Relative intensity distributions along slit S3 aligned on stars #37 and #47. The red line represents the intensity distribution of the He I 2.058 μm emission line. The dashed red line represents the He I 2.113 μm and the green one the H_2 2.121 μm , both are multiplied by a factor 5. The continuum/ $\text{Br}\gamma$ ratio multiplied by a factor 100 is plotted (dashed line). The position of each 1-D spectrum is indicated by a small horizontal segment (solid line). The plot range corresponds to 10.6'' (1pix = 0.05273'').

spectra are represented by horizontal line segments on the plots corresponding to the distribution of the $\text{Br}\gamma$, He I 2.058, 2.113 and H_2 2.121 μm emission lines as well as the continuum emission (Figures 12a, b and c). The length of the segment is proportional to the number of lines integrated along the slit. All the emission lines are continuum subtracted. This figure also indicates the H_2 components C1, C2 and C3 (see Sect. 3.5) of which the distribution intends to clarify the presence of the structures seen in Figure 13 (see Sects. 3.6 and 3.7). From the spectra presented in Figs. 10 and Fig. 11 we derive a ratio He I 2.113 μm / $\text{Br}\gamma$ lines

of mean value 0.06 indicating a hot O star of $T_{\text{eff}} \geq 40000$ K (Hanson et al. 2002). Table 6 (col. 8) shows that this ratio is fairly constant across N88A.

In the spectra of stars #37, #41, #42 and #47 (Fig. 10) the He II 2.185 μm absorption is not detected, if present. The detection is difficult because of our low signal/noise and our low spectral resolution. The NIII 2.115 μm is not detected either. When the He II 2.185 μm absorption line is not present (Bik et al. 2005), the spectral type of a star should be later than O8 V, which is the case for our four resolved stars. As seen above, the spectral type of the ionizing source of the whole nebula derived from radio and $\text{H}\beta$ flux ranges between O4 and O6. We will adopt a type O5 for our computation. Its comparison with the type O8 V derived from our spectroscopy for the bright central star #2-41 clearly shows that other massive stars must contribute to the ionization of N88A. The flux excess between the ionizing star #41 of type O8 V and the O5 type derived from the flux could be produced by at least five O8 V stars. The massive stars #37, #42, and #47 (Fig. 4) located in the upper part of the CM diagram could be good candidates for the ionization of N88A. The 3cm radio peak centered at $(\alpha, \delta) = (1^{\text{h}}24^{\text{m}}7^{\text{s}}.9, -73^{\circ}9'4'')$ and our images show that the radio peak coincides perfectly with the central bright star #41 ($1^{\text{h}}24^{\text{m}}7^{\text{s}}.95, -73^{\circ}9'3''8$) (Table 4). This strong radio emission superposing the NIR emission $\text{Br}\gamma$ line 2.16 μm line (Figs. 12a and b) is characteristic for an H II region.

3.5.2. The N88A-cl cluster

On the JHK image of Lada et al. (2000) the Trapezium region of size of $\sim 0.75 \times 0.75$ pc located at a distance of 450 pc, contains four bright central massive stars and a plethora of low-mass stars with IR-excess. In our Ks-band N88A has approximately a similar diameter (Fig. 4b) and contains also the cluster N88A-cl. This cluster contains other resolved stars not identified by DAOPHOT as well as unresolved stars in crowded groups (Fig. 4b). Among these stars the four brightest ones analyzed using the JHK bands also exhibit IR-excess. N88A with its cluster appears morphologically comparable with the Trapezium region, and other compact star formation regions of similar size, like SH2 269 in our galaxy, and N159-5 in the LMC (Testor et al. 2007). N88A-cl can also be compared with the pre-main-sequence (PMS) clusters, candidate YSOs, of size 0.24 pc to 2.4 pc found in SMC-N66 (Gouliermis et al. 2008). Like N88A-cl these PMS clusters are found coinciding with [OIII], $\text{H}\alpha$ and H_2 emission peaks (see Sect. 3.7). Their clustering properties are similar to the star forming region Orion despite its higher metallicity (Hennekemper et al. 2008).

3.5.3. The bright central star #41

In the Trapezium, the four ionizing bright stars lie within a diameter of ~ 0.05 pc. At the distance of the SMC our spatial resolution is ~ 0.03 pc (~ 6900 AU). With this relative lower spatial resolution it is not excluded that the ionizing star #41 could also be a tight young cluster. This assumption is strengthened by the photometry. The magnitude of this star obtained using the PSF is 15.05 (Table 4). Its dereddened magnitude derived with $A_K = 0.58$ mag corresponds to a mass of $\sim 40 M_{\odot}$ (Fig. 8). Using the parameters for O stars of Vacca (1996), we classify #41 as a




Fig. 13. The H_2 2.121 μm emission (continuum subtracted) showing the three components C1, C2 and C3 of the shell (a) as well as the larger shell (b). The R.A. and Dec. offsets (arcsec) are from star #41 represented by a cross (coordinates 0,0). The image size is $42'' \times 40''$, corresponding to $\sim 13 \times 12$ pc.

O6.5 V type star instead of O8 V derived by spectroscopy, and it could also be multiple.

3.6. The continuum dust emission

Through the JHK-band, N88A presents a relatively bright nebular continuum emission centered on star #41 (Figs. 4a and 5). In the L'-band the continuum is less homogeneous, due to the brightness of L1, L2, L3 and L4. Although the signal to noise ratio is not high, L4 appears very faint in the y-band image (Fig. 4c). The nature of this continuum is not clear. The relatively strong L-band excess of $K-L' = 2$ mag derived from the integration of the whole region N88A (Table 4) supposes that the continuum could come from the emission of circumstellar material (Lada et al. 2000) around resolved and unresolved young stars. These stars could be located mainly at the positions L1, L2, L3 and L4 (Fig. 4d). L4 shows two peaks P1 and P2 visible on the continuum plot of S3 (Fig. 12c). However, the continuum emission could also be formed by interstellar dust associated with the gas of the CO cloud (Testor et al. 1985, HM99, Stanamirovic et al. 2000, Israel et al. 2003). In Fig. 4c the strong optical emission at the position of the 'dark spot' could be explained by strong dust scattering reflecting the light of at least star #37. The nature of the con-

tinuum emission of N88A should be a combination of the two possibilities: emission of circumstellar material and/or dust associated with the gas. Along the slits S1, S2 and S3 the intensity distribution of the continuum near the $\text{Br}\gamma$ line shows a strong continuum/ $\text{Br}\gamma$ ratio of 0.30-0.4 over a range of $6''$ around star #41 (Figs. 12ab). In these figures the broadness of the continuum and $\text{Br}\gamma$ distribution are similar (FWHM $\sim 1.8''$).

3.7. H_2 emission

In Testor et al. (2005) the profile of the H_2 emission along the slit corresponding to our spectrum S1 appears in the form of two blended profiles, due to the low spatial resolution spectroscopy. Thanks to the high spatial resolution of our new data, the complex morphology of the H_2 emission in N88A is revealed both by imagery and spectroscopy. Fig. 13 shows a bidimensional image of the H_2 emission ($v=1-0$ S(1) line). This image is achieved by subtracting the image in the 2.24 μm filter, which allows the passage only of the continuum radiation from the image in the 2.13 μm filter. In this H_2 image, N88A resembles a circular shell (a) of diameter $\sim 3''$ with three maxima labelled C1, C2 and C3 (Fig. 13). Within (a) there is a cavity suggesting radiation pressure, especially from the four central stars. The

Table 6. Relative integrated HeI, HI and H₂ emission-line fluxes in N88A for 14 positions extracted across S1, S2 and S3 (Fig. 12abc).

Id	HeI 2.058 μ m	HeI 2.113 μ m	H ₂ 1-0 S(1) 2.121 μ m	Br γ 2.166 μ m	H ₂ 1-0 S(0) 2.223 μ m	H ₂ 2-1 S(1) 2.247 μ m	I(He 2.113 μ m)/I(Br γ)
S1-41	3.18	0.375	0.121	6.07	0.099	0.031	0.062
S1-42	3.36	0.337	0.075	5.94	0.092	0.044	0.057
S2-41	2.38	0.257	0.094	4.76	0.068	0.038	0.054
S3-37	0.76	0.065	0.028	1.21	0.018	0.009	0.054
S3-47	0.60	0.055	0.066	1.01	0.033	0.006	0.054
S1-1	2.43	0.24	0.199	4.30	0.126	0.069	0.054
S1-2	0.17	0.017	0.094	0.31	0.054	0.035	0.057
S2-1	1.49	0.16	0.150	3.24	0.093	0.048	0.049
S3-1	0.67	0.065	0.049	1.11	0.033	0.009	0.059
S3-2	0.58	0.057	0.026	0.95	0.018	0.009	0.060
S3-3	0.02	0.002	0.017	0.04	0.010	0.008	0.056
S3-4	0.02		0.019	0.02	0.009	0.009	
S3-p1	0.74	0.092	0.020	1.42	0.014	0.009	0.065
S3-p2	0.89	0.090	0.036	1.55	0.034	0.008	0.058

structure C2 is very bright and extended along the direction northeast-southwest and coincides with the ionization front detected by HM99. C2 has a sharp extension in the direction of star #47. The H₂-band image also shows that the bow-shaped filament located east of N88A seems to belong to a second shell (b) of diameter 7'' centered on N88B (Fig. 13). The shells (a) and (b) seem to be in interaction approximatively at the position C1.

Unlike the spectra obtained with ISAAC (Testor et al. 2005) the high-spatial resolution long-slit spectra S1, S2 and S3 allow to resolve the inner structures and stars of N88A. In the direction of the slit S1, the two H₂ emission structures C1 and C2 are well resolved (Fig. 12a) and separated by $\sim 2''$. C1 distant of $\sim 0.5''$ from #41 coincides with the absorption lane observed in optical images by Kurt et al. (1999) and HM99. In the direction of the slit S2, only the structure C1 is seen east of star #41 (Fig. 12b). In the direction of the slit S3, the well seen structures C2 and C3 (Fig. 12c) coincide with the stars #47 and #37 respectively. According to Rubio et al. (2000) massive star formation could be taking place in dense H₂ knots associated with molecular clumps. According to Gouliermis et al. (2008) PMS clusters could be candidate YSOs. These results strengthen our assumption that N88A-cl could contain YSOs.

From Israel & Koorneef (1988), the molecular hydrogen emission may be caused either by shock excitation due to embedded stars, or by fluorescence of molecular material in the ultraviolet radiation field of the OB stars exciting the H II region in the molecular cloud. They conclude that in the MCs, shock excitation of H₂ is only expected very close to (i.e. 0.15pc) stars embedded in a molecular cloud. At a larger distance, radiative excitation of H₂ by the UV radiation field of the OB stars is the only mechanism. Their spectrophotometry with a large aperture (10'') made difficult a precise determination of the brightness of the lines 2-1(S1), 1-0(S1) and 1-0(S0) usually considered between shocks and radiative excitation. These lines deblended when necessary have been derived from our low-spectral-resolution spectra S1, S2 and S3 crossing the zones C1, C2 and C3 as well as the stars #41, #42, #37 and #47 and their intensities are shown in Table 6. None of these lines suffers from atmospheric absorption, considering

a V_{lsr} of 147 km s⁻¹ (Israel et al. 2003), as it can be derived from the solar spectrum atlas (Livingston & Wallace 1991) with the help of a useful home-made software¹. The lines may suffer from differential reddening. Mathis (1990) estimates that the effect for Galactic Sources follows a power-law in the J-, H-, and K-bands: $I_1/I_2 = (\lambda_1/\lambda_2)^{-1.7}$. The effect on the v=1-0 S(0) and v=2-1 S(1) lines would be that they are overestimated by 10%. However, this is based entirely on Galactic Sources. To the best of our knowledge, a differential reddening law has not been determined for Extra-Galactic Sources. Moreover, since the effect is already within our observational uncertainty, we choose to ignore it. In Figure 14 the strengths of the 2.247 μ m 2-1(S1) and 2.223 μ m 1-0(S1) lines are shown relative to the 2.121 μ m 1-0(S1) line for all the objects in Table 6. For radiative excitation (PDRs), the usual criteria are that these ratios should range from 0.5 to 0.6 and from 0.4 to 0.7 respectively (Black & van Dishoeck 1987), while for shock-excitation with T=2000K they should be 0.08 and 0.21 respectively (Shull & Hollenbach 1978). We show in Figure 14 the results from Draine & Bertoldi (1996) reported by Hanson et al. (2002) for high density ($n_H=10^6$ and $\chi=10^4$ and 10^5) and low density ($n_H=10^4$ and $\chi=10^2$) PDRs. We also show in this figure results for more recent and elaborated PDR models (Le Petit et al. 2006) [for n_H ranging from 10^4 to 10^7 and χ from 10^3 to 10^7] as well as for shock models (Flower & Pineau des Forêts 2003) [for the same range of n_H , v_S from 10 to 60 km s⁻¹, the magnetic scaling factor b from 0 to 10 and an ortho/para ratio of 3]. None of these models, either PDRs or shocks really fit with our observations, with the exception of the objects S3-p1 and p2 as well as S3-1, well inside the nebulosity, which could fit with shocks (relatively high velocity v_S , 30 to 50 km s⁻¹) and low b ~ 0.1 . Nevertheless, apart from the three H₂ lines mentioned above, four additional ones are observed: 2-1 S(2), 2-1 S(3), 3-2 S(1) and especially 3-2 S(2) at respectively 2.154, 2.073, 2.386 and 2.286 μ m. These lines are more sensitive to absorption by atmospheric lines, depending in fact on the accuracy of the v_{lsr} . The first one is the less affected by positive or negative velocity shift, whilst the

¹ <http://www.u-cergy.fr/LERMA-LAMAP/informatique/raiesH2/index.html>




Fig. 14. 2-1 S(1)/1-0 (S1) vs. 1-0 S(0)/1-0 S(1) ratio at different positions in the HEB N88A.

two following are slightly absorbed up to 180 km s^{-1} but may suffer a 50 % absorption at 140 km s^{-1} . The last one is free of absorption between 115 and 150 km s^{-1} and does not appear to be blended with any other lines. Possible turbulence in the emitting region may broaden the lines, then lowering the effect due to atmospheric absorption. In any case the concomitant appearance of lines emanating from high v or J as the 3-2 S(2) H_2 line shows a clear trend in favor of the major presence of PDR excitation for most of the observed objects, without nevertheless totally excluding the additional presence of shock excitation. Clearly higher, both spatial and spectral resolutions are required to progress in the knowledge of these objects.

4. Conclusions

We present high spatial resolution imaging of FWHM $\sim 0.12'' - 0.25''$ in the JHKL'-band of the HEB N88A and its immediate environment, and the main results are as follows:

N88A is associated with a cluster that contains at least thirteen stars centered approximatively on the bright central star #41, that could be multiple.

N88A coincides perfectly with the 3cm radio peak and should be ionized not only by the star #41 classified of type O8 V, but also by other low to high-mass stars.

From analysis of the JHK CC diagram we found four possible MYSO star candidates in the N88A cluster, as well as three probable YSOs in the red bow east of N88A. In N88 at least 30% of the detected stars have an IR-excess.

From the K-L excess we found that the core L1-C in N88A should be a heavily embedded, high mass protostar of Class I.

The continuum emission at the position of #41 is very bright and represents about 30% of the $\text{Br}\gamma$ emission peak.

The H_2 infrared emission in N88A resembles a shell formed mainly by three peaks of which one coincides with the ionization front.

We show that the excitation mechanism may be caused predominantly by PDRs, without excluding combination with shocks.

The morphology of N88A could be comparable with galactic regions such as the nearby Trapezium region in the Orion nebula.

Future JHK band imaging data, using higher spatial resolution and longer wavelengths, as L' and M' provided by the NACO S13 camera are still needed to disentangle the IR-excess origin in N88A. Higher spectral resolution spectra are also required to obtain a better analysis of the different spectral lines like the $\text{Br}\gamma$ emission. These new observations should allow to investigate more thoroughly the HEB N88A. This object, which is the brightest, the most excited and reddened of the MCs, presents a unique opportunity to progress in the knowledge of newborn massive stars in regions of low metallicity.

Acknowledgements. We would like to thank the Directors and Staff of the ESO-VLT for making possible these observations and particularly the NACO team for their excellent support. JLL, LK and SD would like to acknowledge the support of the French PCMI program "Physico Chimie du Milieu Interstellaire", funded by the CNRS. This research has made use of the Simbad database, VizieR and Aladin operated at CDS, Strasbourg, France, and the NASA's Astrophysics Data System Abstract Service.

References

- Bik, A., Kaper, L., Hanson, M.M., Smits, M. 2005, A&A, 440, 121
- Bik, A., Kaper, Waters, L.B.F.M. 2006, A&A, 455, 561
- Black, J.H., van Dishoeck, E.F. 1987, ApJ, 322, 412

- Cesaroni, R., Galli, D., Lodato, G., Walmsley, C.M., Zhang, Q. 2007, in *Protostars and Planets V*, ed B. Reipurth, D. Jewitt, & K. Keil, 197
- Charmandaris, V., Heydari-Malayeri, M., Chatzopoulos, E. 2008, *A&A*, 487, 567
- Chen, C.H.R., Chu, Y.-H., Gruendl, R. A., Gordon, K.D., Heisch, F. 2009, *ApJ*, 695, 511
- Cioni, M.-R., Loup, C., Habing, H.J., et al. 2000, *A&AS*, 144, 235
- Cole, A. 1998, *ApJ*, 500, L137
- Cutri, R. M., et al. 2003, *The IRSA 2MASS All-Sky Point Source Catalog*, NASA/IPAC Infrared Science Archive. <http://irsa.ipac.caltech.edu/applications/Gator/>,
- Draine, B.T., Bertoldi, F., 1996, *ApJ*, 468, 269D
- Flower, D. R., Pineau des Forêts, G. 2003, *MNRAS*, 343, 390
- Gouliermis, D., Chu, Y.-H., Henning, T., Brandner, W., Gruendl, R. 2008, *ApJ*, 688, 1050
- Hanson, M., Luhman, K., Rieke. 2002, *ApJS*, 138, 35
- Henize, K. G. 1956, *ApJS*, 2, 315
- Hennekemper, E., Gouliermis, D., Henning, T., Brandner, W., Dolphin, A. 2008, *ApJ*, 672, 914
- Heydari-Malayeri, M., & Testor, G. 1982, *A&A*, 111, L11
- Heydari-Malayeri, M., Le Bertre, T., Magain, P. 1988, *A&A*, 195, 230
- Heydari-Malayeri, M., Charmandris, V., Deharveng, L., Rosa, M.R., Zinnecker, H. 1999, *A&A*, 347, 841
- Heydari-Malayeri, Rosa, M.R., M., Charmandris, V., Deharveng, Martins, F., Meynadier, F., Shaerer, D., Zinnecker, H. 2007, *arXiv0707.1209H*
- Hodge P.W., Wright, F.W. 1977, *The Small Magellanic Cloud*. University of Washington Press, Seattle
- Indebetouw, R., Johnson, K., Conti, P. 2004, *AJ*, 128, 2206
- Israel, F.P., Koorneef, J. 1988, *A&A*, 190, 21
- Israel, F.P., Koorneef, J. 1991, *A&A*, 248, 404
- Israel, F.P., Johansson, L., Rubio, M., et al. 2003, *A&A*, 406, 817
- Kovacs, G. 2000, *A&A* 360, L1
- Kurt,C., Dufour, R., Garnett, D., et al. 1999, *ApJ*, 518, 246,261
- Lada, C.J., A.A Muench and five authors 2000, *ApJ*, 120,3162
- Livingston, W., Wallace, L. 1991, *NSO Technical Report*, Tucson: National Solar Observatory, National Optical Astronomy Observatory
- Lejeune, T., Shaerer, D. 2001, *A&A*, 366, 538
- Le Petit, F., Nehmé, C., Le Bourlot, J. and Roueff, E. 2006, *ApJS*, 164, 506
- Mathis, J. 1990, *ARA&A*, 28, 37
- Maercker, M., Burton, G. 2005, *A&A*, 438, 663
- Martin-Hernandez, N.L., Peeters, E., Tielens, A.G.G.M 2008, *A&A*, 489, 1189
- McGee, R.X., Newton, L.M. 1982, *Proc. Astron. Soc. Australia*, 4, 308
- McNamara, D. H., Feltz, K. A. 1980, *PASP*, 92, 587
- Okano, K. <http://www.asahi-net.or.jp/~rt6k-okn/its98/ddp1.htm>
- Pugliese, G., Christou, J., Koehler, R., Drummond, J. 2002, *AAS* 201, 5505
- Rieke, G., Lebofsky, M. 1985, *ApJ*, 288, 618
- Rubio, M., Contursi, A., Lequeux, J., Prost, R., Barba, R., Boulanger, F., Cesarsky, D., Maoli, R. 2000, *A&A*, 359, 1139
- Schmeja, S., Gouliermis, D., Klessen, R. 2009, *ApJ*, 694, 367
- Shull, J. M & Hollenbach, D. J., 1978, *ApJ*, 220, 525
- Simon, J.D., Bollato, A., Stanimirovic, S., Sha, R., Leroy, A., Sandstrom, K. 2006, *astro-ph/0603834v1*
- Smith, L. J., Noriss, R.P. F., Crowther, P. A. 2002, *MNRAS*, 337, 1309
- Stanimirovic, S., Staveley-Smith, L., van der Hulst, J.M., Bontekoe, T.J.R., Kester, D.J.M., Jones, P.A. 2000, *MNRAS*, 315, 791
- Stanghellini, L., Villaver, E., Shaw, R., Mutchler, M. 2003, *ApJ*, 598, 1000
- Stetson, P. B. 1987, *PASP*, 99, 191
- Tapia, M., Persi, P., Roth, M., Marenzi, A., Testi, L., Vanzi, L. 2003, *MNRAS*, 339, 44
- Testor, G., Pakull, M. 1985, *A&A*, 145, 170
- Testor, G. 2001, *A&A*, 372, 667
- Testor, G., Lemaire, J.L., Field, D. 2003, *A&A*, 407, 905
- Testor, G., Lemaire, J.L., Field, D., Callejo, G. 2005, *A&A*, 434, 497
- Testor, G., Lemaire, J.L., Field, D., Kristensen, L.E., Diana, S. 2007, *A&A*, 469, 459
- Vacca, W.D., Garmany, C.D., & Shull, J.M. 1996, *ApJ*, 460, 914
- Westerlund, B.E. 1997, *The Magellanic Clouds* (Cambridge: Cambridge Univ.Press), pp. 28-34
- Whitney, B. A., Indebetouw, R., Bjorkman, J.E., Wood, K. 2004, *ApJ*, 617, 1177
- Wilcots, E.M. 1994b, *AJ*, 108, 1674
- Zaritsky, D., Harris, J., Thompson, I., Grebel, E., K., Massey, P. 2002, *AJ*, 123, 855